

AD-A039 208

INDIANA UNIV BLOOMINGTON DEPT OF MATHEMATICS
ASYMPTOTIC NORMALITY AND CONVERGENCE RATES OF LINEAR RANK STATISTICS (U)
1977 M L PURI, N S RAJARAM

F/G 12/1

AF-AFOSR-2927-76

UNCLASSIFIED

AFOSR-TR-77-0451

NL

1 OF 1
AD-A039 208

12/1

END

DATE
FILMED
6-77

Asymptotic Normality and Convergence Rates
of Linear Rank Statistics Under AlternativesBy Madan L. Puri and Navaratna S. Rajaram
Indiana University and Kent State University

AD A 039208

Approved for public release;
distribution unlimited.

0. Summary. Asymptotic normality of linear rank statistics under alternatives is proved employing techniques of Chernoff and Savage, (1958). Under suitable assumptions the rate of convergence to normality under alternatives is also obtained for such statistics. The results of the paper are related to those of Hájek (1968) and Hoeffding, (1973). Results on the rate of convergence extend those of Jurečková and Puri, (1976), and Bergström and Puri, (1976).

1. Introduction. Let $\{X_{Ni}, i \geq 1\}$ be a sequence of independent random variables with continuous cdfs (cumulative distribution functions) $\{F_{Ni}, i \geq 1\}$ respectively. Consider a linear rank statistic S_N given by

$$(1.1) \quad S_N = \sum_{i=1}^N c_{Ni} a_N(R_{Ni})$$

where R_{Ni} is the rank of X_{Ni} in (X_{N1}, \dots, X_{NN}) , (c_{N1}, \dots, c_{NN}) are known (regression) constants, and $a_N(1), \dots, a_N(N)$ are "scores" generated by a known real-valued function $\varphi(t)$, $0 < t < 1$ in either of the following ways:

$$(1.2) \quad a_N(i) = \varphi(i/(N+1)), \quad 1 \leq i \leq N$$

$$(1.3) \quad a_N(i) = E \varphi^{(i)}(U_N^{(i)}), \quad 1 \leq i \leq N$$

where $U_N^{(i)}$ is the i th order statistic in a sample of size N from the rectangular distribution over $(0,1)$.

Work supported by the Air Force Office of Scientific Research, A.F.S.C.U.S.A.F., Grant No. AFOSR76-2927. Reproduction in whole or part permitted for any purpose of the United States Government.

AD NO. —
DDC FILE COPY

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH (AFSC)
NOTICE OF TRANSMITTAL TO DDC
This technical report has been reviewed and is
approved for public release IAW AFR 190-12 (7b).
Distribution is unlimited.

A. D. BLOSE
Technical Information Officer

We make the following assumptions:

$$(I A) \quad \max_{1 \leq i \leq N} |c_{Ni}| / s_N^{-1/2} = o(N) \quad (N)$$

$$(I B) \quad |\varphi^{(i)}(t)| \leq K [t(1-t)]^{\delta \cdot i - 1/2}, \quad i = 0, 1; \quad \delta > 0,$$

K a generic constant.

s_N^2 is approximate variance of S_N and is given by (1.7) and (1.8) below.

Our main results are the following:

Theorem 1.1. Let the scores $a_N(i)$, $1 \leq i \leq N$ be defined as in (1.2).
Then, under the assumptions (I A) and (I B),

$$(1.4) \quad \sup_x |P\left(\frac{S_N - \mu_N}{s_N} \leq x\right) - \Phi(x)| \rightarrow 0 \quad \text{as } N \rightarrow \infty, \quad s_N \neq 0$$

where

$$\Phi(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^x e^{-t^2/2} dt,$$

$$(1.5) \quad \mu_N = \sum_{i=1}^N c_{Ni} \int_{-\infty}^{\infty} \varphi(H(x)) dF_{Ni}(x),$$

$$(1.6) \quad H(x) = \frac{1}{N} \sum_{i=1}^N F_{Ni}(x),$$

$$(1.7) \quad s_N^2 = \sum_{i=1}^N s_{Ni}^2; \quad s_{Ni}^2 = \text{Var}(A_{Ni}(X_{Ni})),$$

$$(1.8) \quad A_{Ni}(x) = \frac{1}{N} \sum_{j=1}^N (c_{Ni} - c_{Nj}) \int \{I_{[x \leq y]} - F_{Ni}(y)\} \varphi'(H(y)) dF_N(y),$$

and

$$I_{[x \leq y]} = \begin{cases} 0 & \text{if } y < x \\ 1 & \text{if } y \geq x \end{cases}$$



Corollary 1.1. Let $\varphi(t) = F^{-1}(t)$, where F is a cdf. Let the scores $a_N(i)$, $1 \leq i \leq N$ be given by (1.3). Then, under the assumptions (IA) and (IB), the conclusions of Theorem 1.1 hold.

Theorem 1.2. Let the scores $a_N(i)$, $1 \leq i \leq N$ be given by (1.2) and the assumption (IA) be satisfied. Let

$$(1.10) \quad \sup_x |\varphi'(x)| = \|\varphi'\| < \infty$$

then

$$(1.11) \quad \sup_x \left| P\left(\frac{S_N - \mu_N}{s_N} \leq x\right) - \Phi(x) \right| \leq c_1 \ell_{3,N} + \Delta_N$$

where

$$(1.12) \quad \ell_{3,N} = \rho_N^3 / s_N^3, \quad c_1 = 0.7985$$

$$(1.13) \quad \rho_N^3 = \sum_{i=1}^N \rho_{Ni}^3, \quad \rho_{Ni}^3 = E |A_{Ni}(X_{Ni})|^3, \text{ and } \Delta_N \rightarrow 0$$

Furthermore,

$$(1.14) \quad N^{\frac{1}{2}} \Delta_N = o_p(\|\varphi'\| s_N^{-1} \sum_{i=1}^N |c_{Ni}|)$$

Remark. Theorem 1.1 has been proved by Hájek (1968). His conditions on the score generating function φ are milder than ours but the conclusion of our theorem 1.1 is sharper in the sense that the centering constant μ_N appears naturally in place of ES_N given by Hájek (1968). Corollary 1.1 is an extension of a similar result proved by Chernoff and Savage (1958) for the two sample problem and serves to remove some of the complications encountered in Hoeffding (1973). Theorem 1.2 is related to the results of Bickel (1972), Jurečková-Puri (1975), and Bergström-Puri (1976). However, the bounds obtained in these papers are non-random and thus sharper than ours. On the other hand, our conditions on the score generating function φ are milder. We believe that the theorem 1.2 is true even when the condition (1.10) is replaced by the assumption (IB). At the present time, the theory of asymptotic expansion for sums of dependent random variables is still at a rudimentary stage (Stein (1970)) and it is doubtful if the random term Δ_N can be removed without additional assumptions on the underlying distributions.

Asymptotic Normality of S_N . The proof of theorem 1.1 (and corollary 1.2) will be along the lines of the Chernoff-Savage theorem (1958) as given in Puri and Sen (1971) with some modifications necessitated by greater generality of the present problem.

First we introduce the following notations:

$$(2.1) \quad H_N(x) = \frac{1}{N} \sum_{i=1}^N I_{[X_{Ni} \leq x]}$$

$$(2.2) \quad H(x) = \frac{1}{N} \sum_{i=1}^N F_{Ni}(x)$$

$$(2.3) \quad C_N(x) = \sum_{i=1}^N C_{Ni} I_{[X_{Ni} \leq x]}$$

$$(2.4) \quad C(x) = \sum_{i=1}^N C_{Ni} F_{Ni}(x) .$$

Note that the functions $H_N(x)$ and $C_N(x)$ are stochastic variables whereas $H(x)$ and $C(x)$ are non-random though depending on N .

Then the following inequalities are obvious:

$$(2.5) \quad |C_N(x)| \leq N \max_{1 \leq i \leq N} |C_{Ni}| H_N(x)$$

$$(2.6) \quad |C(x)| \leq N \max_{1 \leq i \leq N} |C_{Ni}| H(x), \quad -\infty < x < \infty .$$

Proof of Theorem 1.1. We rewrite S_N defined in (1.1) as

$$(2.7) \quad \begin{aligned} S_N &= \int_{-\infty}^{\infty} \varphi\left(\frac{N}{N+1} H_N(x)\right) dC_N(x) \\ &= \mu_N + B_{1N} + B_{2N} + \sum_{i=1}^3 D_{iN} \end{aligned}$$

where μ_N is given by (1.5), and

$$(2.8) \quad B_{1N} = \int_{-\infty}^{\infty} \varphi(H(x)) d(C_N(x) - C(x)) ,$$

$$(2.9) \quad B_{2N} = \int_{-\infty}^{\infty} (H_N(x) - H(x)) \varphi'(H(x)) dC(x) ,$$

$$(2.10) \quad D_{1N} = \frac{-1}{N+1} \int_{-\infty}^{\infty} H_N(x) \varphi'(H(x)) dC_N(x) ,$$

$$(2.11) \quad D_{2N} = \int_{-\infty}^{\infty} (H_N(x) - H(x)) \varphi'(H(x)) d(C_N(x) - C(x)) ,$$

$$(2.12) \quad D_{3N} = \int_{-\infty}^{\infty} \left\{ \varphi\left(\frac{N}{N+1} H_N(x)\right) - \varphi(H(x)) - \left(\frac{N}{N+1} H_N(x) - H(x)\right) \varphi'(H(x)) \right\} dC_N(x) .$$

The proof will be accomplished if we establish the following:

$$(a) \quad |\mu_N| < \infty ,$$

$$(b) \quad (B_{1N} + B_{2N})/s_N \text{ is asymptotically normal,}$$

$$(c) \quad D_{iN} = o_p(s_N), \quad i=1,2,3 .$$

Proof of (a). Using (2.4) and (2.6), we obtain

$$(2.13) \quad |\mu_N| \leq N \max_{1 \leq i \leq N} |C_{Ni}| \int_{-\infty}^{\infty} |\varphi(H(x))| dH(x) < \infty, \text{ by assumption (I B) .}$$

Proof of (b). To prove (b), we shall verify the Liapunov condition for B_{1N}/s_n and B_{2N}/s_n . Integrating B_{2N} by parts, we obtain

$$(2.14) \quad B_{2N} = [H_N(x) - H(x)] B^*(x) \Big|_{-\infty}^{\infty} - \int_{-\infty}^{\infty} B^*(x) d[H_N(x) - H(x)]$$

where

$$B^*(x) = \int_{x_0}^x \varphi'(H(x)) dC(x)$$

where x_0 is determined arbitrarily such that $H(x_0) > 0$.

We first consider $\int_{-\infty}^{\infty} B^*(x) d[H_N(x) - H(x)]$

$$(2.15) \quad \int_{-\infty}^{\infty} B^*(x) d[H_N(x) - H(x)] = \sum_{i=1}^N \left[B^*(x_{Ni}) - E B^*(x_{Ni}) \right] / N$$

We verify the condition $\frac{1}{s_N^{2+\delta'}} \sum_{i=1}^N E \left[|B^*(x_{Ni}) - E B^*(x_{Ni})| / N \right]^{2+\delta'} \rightarrow 0$

as $N \rightarrow \infty$ for some $\delta' > 0$. To verify this, it suffices to show that

$$(2.16) \quad \frac{1}{s_N^{2+\delta'}} \sum_{i=1}^N E \left[|B^*(x_{Ni})| / N \right]^{2+\delta'} \rightarrow 0, \text{ as } N \rightarrow \infty.$$

Choose $\delta' > 0$ such that $(2 + \delta')(\delta - \frac{1}{2}) > 1$. Then

$$(2.17) \quad \frac{1}{s_N^{2+\delta'}} \sum_{i=1}^N E |B^*(x_{Ni}) / N|^{2+\delta'} = \frac{1}{s_N^{2+\delta'} \cdot N^{2+\delta'}} \sum_{i=1}^N \int_{-\infty}^{\infty} \left| \int_{x_0}^x \varphi'(H(y)) dC(y) \right|^{2+\delta'} dF_{Ni}(x)$$

$$\leq \frac{1}{s_N^{2+\delta'} \cdot N^{2+\delta'}} \sum_{i=1}^N N^{2+\delta'} \max_{1 \leq i \leq N} |C_{Ni}|^{2+\delta'} \int_{-\infty}^{\infty} \left| \int_{x_0}^x \varphi'(H(y)) dH(y) \right|^{2+\delta'} dF_{Ni}(x)$$

$$\leq \left\{ \max_{1 \leq i \leq N} |C_{Ni}| / s_N \right\}^{2+\delta'} \sum_{i=1}^N \int_{-\infty}^{\infty} \left\{ |\varphi(H(x))| + |\varphi(H(x_0))| \right\}^{2+\delta'} dF_{Ni}(x)$$

$$= O(N^{-\delta'/2}) \int_{-\infty}^{\infty} \left\{ |\varphi(H(x))| + |\varphi(H(x_0))| \right\}^{2+\delta'} dH(x).$$

We now show that

$$\frac{\beta(x)}{s_N} \Big|_{-\infty}^{\infty} = o_p(1), \text{ where } \beta(x) = [H_N(x) - H(x)] B^*(x) \quad (\text{see (2.14)}).$$

We note that

$$\begin{aligned}
\left| \frac{f_N(x)}{s_N} \right| &= \frac{1}{N} |H_N(x) - H(x)| \cdot \frac{1}{s_N} \left| \int_{x_0}^x \varphi'(H(y)) dC(y) \right| \\
&\leq \frac{1}{N} |H_N(x) - H(x)| O(1) \left| \int_{x_0}^x \varphi'(H(y)) dH(y) \right| \\
&\leq k \frac{1}{N} |H_N(x) - H(x)| \{H(x)(1-H(x))\}^{\delta-1/2}
\end{aligned}$$

Now since $\forall \epsilon > 0, \delta' > 0, \exists c(\epsilon, \delta') \ni$

$$(2.18) \quad P \left[\sup_x \frac{1}{N} \frac{|H_N(x) - H(x)|}{\{H(x)(1-H(x))\}^{\delta-1/2}} > c(\epsilon, \delta') \right] < \epsilon,$$

it follows that with probability $> 1 - \epsilon$,

$$\left| \frac{f_N(x)}{s_N} \right| \leq k \{H(x)(1-H(x))\}^{\delta-\delta'} c(\epsilon, \delta') \rightarrow 0 \text{ as } x \rightarrow \pm \infty$$

by choosing $\delta' < \delta$.

Thus the Liapunov condition (2.16) is satisfied for B_{2N}/s_N . The verification (of the Liapunov condition) for B_{1N}/s_N is similar, and the same is true for $(B_{1N}+B_{2N})/s_N$ by using the C_r -inequality. This proves (b).

Proof of (c).

$$(2.19) \quad \left| \frac{1_N}{N} \right| \leq \frac{1}{Ns_N} \left| \sum_{i=1}^N \varphi'(H(X_{Ni})) C_{Ni} \right| \leq \frac{1}{N} \sum_{i=1}^N V_{Ni}$$

where

$$(2.20) \quad V_{Ni} = \left| \varphi'(H(X_{Ni})) \frac{C_{Ni}}{s_N} \right|$$

To establish that $\frac{1}{N} \sum_{i=1}^N V_{Ni} \rightarrow 0$ in probability, it suffices to show that

$$(2.21) \quad \frac{1}{N} \sum_{i=1}^N E|V_{Ni}|^\alpha < \infty \text{ for some } 0 < \alpha < 1 \text{ (cf. Loève (1963), page 241).}$$

Taking $\alpha = 2/3$,

$$\begin{aligned} \frac{1}{N} \sum_{i=1}^N E |V_{Ni}|^{2/3} &\leq \frac{K}{N} \max_{1 \leq i \leq N} \left| \frac{C_{Ni}}{s_N} \right|^{2/3} \times \\ &\times \sum_{i=1}^N \int_{-\infty}^{\infty} \{H(x)(1-H(x))\}^{\frac{2\delta}{3}-1} dF_{Ni}(x) \\ &\leq K \int_0^1 \{u(1-u)\}^{\frac{2\delta}{3}-1} du < \infty \text{ uniformly in } N. \end{aligned}$$

Now consider

$$D_{2N} = \int_{-\infty}^{\infty} (H_N(x) - H(x)) \varphi'(H(x)) d(C_N(x) - C(x))$$

Noting (2.18), it follows that with probability $> 1-\epsilon$,

$$|H_N(x) - H(x)| |\varphi'(H(x))| \leq \frac{K}{N^2} C(\epsilon, \delta') \{H(x)(1-H(x))\}^{\delta-\delta'-1}$$

Setting $0 < \delta^* = \delta - \delta'$ Choosing $\delta' < \delta$, it suffices to show that

$$\begin{aligned} (2.22) \quad \frac{1}{s_N N^2} \int_{-\infty}^{\infty} \{H(x)(1-H(x))\}^{\delta^*-1} dC_N(x) &= \frac{1}{s_N N^2} \sum_{i=1}^N C_{Ni} \times \\ &\times \{H(X_{Ni})(1-H(X_{Ni}))\}^{\delta^*-1} \end{aligned}$$

Setting

$$(2.23) \quad V_{Ni} = \frac{\frac{1}{2} C_{Ni}}{s_N} \{H(X_{Ni})(1-H(X_{Ni}))\}^{\delta^*-1},$$

we have to show that

$$(2.24) \quad \frac{1}{N} \sum_{i=1}^N \{V_{Ni} - E V_{Ni}\} \rightarrow 0 \text{ in probability.}$$

This will follow if we show that for some $\alpha > 0$,

$$(2.25) \quad N^{-(1+\alpha)} \sum_{i=1}^N E |V_{Ni}|^{1+\alpha} \rightarrow 0$$

Choose $\alpha > 0$ such that $(1+\alpha)(\delta^* - 1) > -1$ (i.e. $0 < \alpha < \frac{\delta^*}{1-\delta^*}$).
Then

$$\begin{aligned}
 (2.26) \quad & N^{-(1+\alpha)} \sum_{i=1}^N E |V_{Ni}|^{1+\alpha} \leq \frac{1}{\alpha N \cdot N} \left| \frac{N^{\frac{1}{2}} \max_{1 \leq i \leq N} |C_{Ni}|}{s_N} \right|^{1+\alpha} \times \\
 & \times \sum_{i=1}^N E \{H(X_{Ni})(1-H(X_{Ni}))\}^{(1+\alpha)(\delta^*-1)} \\
 & = O(1) \frac{1}{N^\alpha} \int_{-\infty}^{\infty} \{H(x)(1-H(x))\}^{(1+\alpha)(\delta^*-1)} dH(x) \rightarrow 0, \text{ as } N \rightarrow \infty.
 \end{aligned}$$

This proves (2.25).

Finally, consider

$$D_{3N} = \int_{-\infty}^{\infty} \left\{ \varphi\left(\frac{N}{N+1} H_N(x)\right) - \varphi(H(x)) - \left(\frac{N}{N+1} H_N(x) - H(x)\right) \varphi'(H(x)) \right\} dC_N(x)$$

We have to show that $\frac{D_{3N}}{s_N} = o_p(1)$. Denote

$$(2.27) \quad C_{3N} = \frac{D_{3N}}{\frac{1}{2} s_N},$$

and note that

$$(2.28) \quad |C_{3N}| \leq O(1) \int_{-\infty}^{\infty} \left| \varphi\left(\frac{N}{N+1} H_N(x)\right) - \varphi(H(x)) - \left(\frac{N}{N+1} H_N(x) - H(x)\right) \varphi'(H(x)) \right|$$

$$\varphi'(H(x)) |dH_N(x), \text{ since } \max_{1 \leq i \leq N} \frac{|C_{Ni}|}{s_N N^{\frac{1}{2}}} = O(N^{-1}).$$

The proof that the right hand side of (2.28) is $o_p(N^{-\frac{1}{2}})$ follows precisely as in Puri and Sen (1971, pages 401-405). Thus the proof of Theorem 1.1 follows.

Proof of Corollary 1.1. Denote

$$\varphi_N(t) = \sum_{i=1}^{[Nt]} a_N(i), \quad 0 \leq t < 1; \quad a_N(i) = E \varphi(U_N^{(i)}), \quad 1 \leq i \leq N$$

where $[x]$ is the greatest integer $\leq x$; and let

$$(2.29) \quad S_N^* = \int_{-\infty}^{\infty} \varphi_N \left(\frac{N H_N(x)}{N+1} \right) dC_N(x)$$

The proof of the corollary will be accomplished if we show that $(S_N - S_N^*)/s_N \rightarrow 0$ in probability, as $N \rightarrow \infty$.

Lemma 2.1. Under the hypotheses of theorem 1.1 and Corollary 1.1,

$$(2.30) \quad \lim_{N \rightarrow \infty} \varphi_N(t) = \varphi(t)$$

$$(2.31) \quad \left| \int_{-\infty}^{\infty} \left\{ \varphi_N \left(\frac{N}{N+1} H_N(x) \right) - \varphi \left(\frac{N}{N+1} H_N(x) \right) \right\} dC_N(t) \right| = o_p(s_N).$$

Proof. The proof of (2.30) is well known (see Puri and Sen (1971), pages 408-409); and so we prove (2.31).

Using (2.5), we obtain

$$(2.32) \quad \begin{aligned} & \left| \int_{-\infty}^{\infty} \left\{ \varphi_N \left(\frac{N H_N(x)}{N+1} \right) - \varphi \left(\frac{N H_N(x)}{N+1} \right) \right\} dC_N(x) \right| \\ & \leq \max_{1 \leq i \leq N} |C_{Ni}| \sum_{i=1}^N \left| \varphi_N \left(\frac{N H_N(X_{Ni})}{N+1} \right) - \varphi \left(\frac{N H_N(X_{Ni})}{N+1} \right) \right| \\ & = \max_{1 \leq i \leq N} |C_{Ni}| \sum_{i=1}^N \left| \varphi_N \left(\frac{i}{N+1} \right) - \varphi \left(\frac{i}{N+1} \right) \right| \end{aligned}$$

Now by assumption I (A),

$$(2.33) \quad \max_{1 \leq i \leq N} \left| \frac{C_{Ni}}{s_N} \right| \sum_{i=1}^N \left| \varphi_N \left(\frac{i}{N+1} \right) - \varphi \left(\frac{i}{N+1} \right) \right| = O(1) N^{-1/2} \sum_{i=1}^N \left| \varphi_N \left(\frac{i}{N+1} \right) - \varphi \left(\frac{i}{N+1} \right) \right|$$

which $\rightarrow 0$ as $N \rightarrow \infty$ (cf. Puri and Sen (1971), pages 409-411).

This proves the Lemma; and hence the corollary 1.1.

To compute the $\text{var}(B_{1N} + B_{2N})$, note that

$$B_{1N} = \sum_{i=1}^N \varphi(H(X_{Ni})) C_{Ni} + \text{constant}.$$

$$B_{2N} = -\frac{1}{N} \sum_{i=1}^N B^*(X_{Ni}) + \text{constant, where}$$

$$B^*(x) = \int_{x_0}^x \varphi'(H(y)) d\mathcal{C}(y) = \sum_{j=1}^N C_{Nj} \int_{x_0}^x \varphi'(H(y)) dF_{Nj}(y)$$

Noting that $\varphi(H(x)) = \int_{x_0}^x \varphi'(H(y)) dH(y) + \varphi(H(x_0))$, and setting

$$\begin{aligned} A_{Ni}(x) &= \frac{C_{Ni}}{N} \sum_{j=1}^N \int_{x_0}^x \varphi'(H(y)) dF_{Nj}(y) - \frac{1}{N} \sum_{j=1}^N C_{Nj} \int_{x_0}^x \varphi'(H(y)) dF_{Nj}(y) \\ &= \frac{1}{N} \sum_{j=1}^N (C_{Ni} - C_{Nj}) \int_{x_0}^x \varphi'(H(y)) dF_{Nj}(y) \end{aligned}$$

we obtain

$$\text{Var}(B_{1N} + B_{2N}) = \sum_{i=1}^N \text{var}(A_{Ni}(X_{Ni})) .$$

Proof of Theorem 1.2. Using (2.7), we have

$$(2.34) \quad \frac{S_N - u_N}{s_N} = \frac{T_N}{s_N} + \frac{D_N}{s_N}$$

where

$$(2.35) \quad T_N = B_{1N} + B_{2N}, \text{ and } D_N = D_{1N} + D_{2N} + D_{3N} .$$

Denote

$$(2.36) \quad F_N(x) = P\left(\frac{S_N - u_N}{s_N} \leq x\right), \quad G_N(x) = P\left(\frac{T_N}{s_N} \leq x\right)$$

Then

$$\begin{aligned} (2.37) \quad F_N(x) - \Phi(x) &= G_N\left(x - \frac{D_N}{s_N}\right) - \Phi(x) \\ &= \left[G_N\left(x - \frac{D_N}{s_N}\right) - \Phi\left(x - \frac{D_N}{s_N}\right)\right] + \left[\Phi\left(x - \frac{D_N}{s_N}\right) - \Phi(x)\right] \end{aligned}$$

Since (by Polya's theorem), both $F_N(x) - \Phi(x)$ and

$G_N\left(x - \frac{D_N}{s_N}\right) - \Phi\left(x - \frac{D_N}{s_N}\right)$ converge to 0 uniformly in x , it

follows that the random quantity

$$(2.38) \quad \Delta_N = \sup_x \left| \Phi\left(x - \frac{D_N}{s_N}\right) - \Phi(x) \right| \text{ converges to zero.}$$

We now estimate

$$(2.39) \quad \sup_x \left| G_N\left(x - \frac{D_N}{s_N}\right) - \Phi\left(x - \frac{D_N}{s_N}\right) \right| = \sup_x \left| G_N(x) - \Phi(x) \right|$$

Observe that by the Berry-Esseen theorem (cf. Feller (1971), page 544),

$$(2.40) \quad \sup_x \left| G_N(x) - \Phi(x) \right| \leq \frac{C \rho_N^3}{s_N^3}$$

where C can be taken to be .7975 (cf. Van Beek (1972), and Bhattacharya-Ranga Rao (1976)).

From (2.40) and (2.37), we have

$$(2.41) \quad \sup_x \left| F_N(x) - F(x) \right| \leq \frac{C \rho_N^3}{s_N^3} + \Delta_N$$

It remains to show that $N^{\frac{1}{2}} \Delta_N = O_p(\|\varphi'\| s_N^{-1} \sum_{i=1}^N |C_{Ni}|)$

From (2.38), since

$$(2.42) \quad \left| \Phi\left(x - \frac{D_N}{s_N}\right) - \Phi(x) \right| = \left| \frac{D_N}{s_N} \right| \left| \Phi'\left(x - \frac{\alpha D_N}{s_N}\right) \right| \text{ for some } \alpha, 0 \leq \alpha \leq 1$$

$$\leq \frac{1}{\sqrt{2\pi}} \left| \frac{D_N}{s_N} \right|$$

it suffices to show that $N^{\frac{1}{2}} \Delta_N = O_p(\|\varphi'\| s_N^{-1} \sum_{i=1}^N |C_{Ni}|)$.

Now re-arranging the terms of D_N , we obtain

$$(2.43) \quad D_N = \int_{-\infty}^{\infty} \left\{ \varphi\left(\frac{N}{N+1} H_N(x)\right) - \varphi(H(x)) \right\} dC_N(x) - \int_{-\infty}^{\infty} (H_N(x) - H(x)) dC(x)$$

To simplify the proof, we drop the factor $\frac{N}{N+1}$ since it does not affect the conclusion. By the mean-value theorem

$$(2.44) \quad \varphi(H_N(x)) - \varphi(H(x)) = (H_N(x) - H(x)) \varphi'(\xi_N(x)) \text{ for some } \xi_N(x).$$

Hence

$$\left| \frac{N^{\frac{1}{2}}}{s_N} \int_{-\infty}^{\infty} (H_N(x) - H(x)) \varphi'(\xi_N(x)) dC_N(x) \right| \leq \frac{\|\varphi'\|}{s_N} \int_{-\infty}^{\infty} N^{\frac{1}{2}} (H_N(x) - H(x)) dC_N(x)$$

Let $\epsilon > 0$ be given. Then from Puri and Sen (1971), there exists a constant $C(\epsilon)$ such that with probability $> 1 - \epsilon$,

$$\sup_x N^{\frac{1}{2}} |H_N(x) - H(x)| < C(\epsilon)$$

Hence, with probability $> 1 - \epsilon$,

$$(2.45) \quad \left| \frac{N^{\frac{1}{2}}}{s_N} \int_{-\infty}^{\infty} (H_N(x) - H(x)) \varphi'(\xi_N(x)) dC_N(x) \right| \leq \frac{C(\epsilon)}{s_N} \|\varphi'\| \sum_{i=1}^N |C_{Ni}|$$

The proof of

$$(2.46) \quad \left| \frac{1}{s_N} \int_{-\infty}^{\infty} N^{\frac{1}{2}} (H_N(x) - H(x)) \varphi'(H(x)) dC(x) \right| \leq \frac{C(\epsilon)}{s_N} \|\varphi'\| \sum_{i=1}^N |C_{Ni}|$$

in probability is identical.

(2.45) and (2.46) establish the theorem (1.2).

REFERENCES

1. Beek, P. van (1972). An application of the Fourier method to the problem of sharpening the Berry-Esseen inequality. Z. Wahrscheinlichkeits theorie verw. Geb. 23, 187-197.
2. Bergström, H. and Puri, M.L. (1977). Convergence and remainder terms in linear rank statistics. Ann. Statist. 5
3. Bhattacharya, R.N. and Ranga Rao, R. (1976). Normal approximation and asymptotic expansions. John Wiley, New York.
4. Bickel, P.J. (1974). Edge worth expansions in nonparametric statistics. Ann. Statist. 2 , 1-20 .
5. Chernoff, H. and Savage, I.R. (1958). Asymptotic normality and efficiency of certain nonparametric test statistics. Ann. Math. Statist. 29, 972-996.
6. Feller, W. (1971). An introduction to Probability theory and its applications, vol. 2. John Wiley, New York.
7. Hájek, J. (1968). Asymptotic normality of simple linear rank statistics under alternatives. Ann. Math. Statist. 39, 325-346.
8. Hoeffding, W. (1973). On the centering of a simple linear rank statistics. Ann. Statist. 1, 56-66.
9. Jurečková, J. and Puri, M.L. (1975). Order of normal approximation for rank test statistics distribution. Ann. Prob. 3, 526-533.

10. Loève, M. (1963). Probability Theory, 3rd ed. Van Nostrand, New York.
11. Puri, M.L. and Sen, P.K. (1971). Non parametric methods in multivariate analysis. John Wiley, New York.
12. Stein, C. (1970). A bound for error in the normal approximation to the distribution of a sum of dependent random variables. Proc. 6th Berkeley Symp. Math. Stat. and Prob. vol. 2 .

SEARCHED	INDEXED
SERIALIZED	FILED
OCT 1971	
FBI - NEW YORK	
A	

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

17 REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER AFOSR-TR-77-0451	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) ASYMPTOTIC NORMALITY AND CONVERGENCE RATES OF LINEAR RANK STATISTICS UNDER ALTERNATIVES,		5. TYPE OF REPORT & PERIOD COVERED Interim report
7. AUTHOR(s) Madan L. Puri and Navaratna S. Rajaram		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Indiana University Department of Mathematics Bloomington, IN 47401		8. CONTRACT OR GRANT NUMBER(s) VAF-AFOSR-2927-76
11. CONTROLLING OFFICE NAME AND ADDRESS Air Force Office of Scientific Research/NM Bolling AFB, Washington, DC 20332		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 61102F 2304/A5
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) 12 17 p.		12. REPORT DATE 10 1977
		13. NUMBER OF PAGES 15
		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) linear rank statistics; convergence rates; score generating function		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Asymptotic normality of linear rank statistics under alternatives is proved employing techniques of Chernoff and Savage (1958 <u>Ann. Math. Statist.</u> , 29, 972-994). Under suitable assumptions the rate of convergence to normality under alternatives is also obtained for such statistics. The results of the paper are related to those of Hajek (1968, <u>Ann. Math. Statist.</u> , 39, 325-346), and		

402 522-08

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

20 Abstract

Hoeffding (1973, Ann. Statist., 1, 54-66). Results on the rate of convergence extend those of Jurečkova and Puri (1975, Ann. Prob., 526-533), and Bergström and Puri (1977, Ann. Statist.).

UNCLASSIFIED

Indian University
Department of Mathematics
Washington, D.C. 20001

Office of Scientific Research
Johns Hopkins University
Washington, D.C. 20001

UNCLASSIFIED

Approved for public release; distribution unlimited.

Abstracts of papers presented at the 1977 Annual Meeting of the American Statistical Association, held in San Francisco, California, December 12-15, 1977.

UNCLASSIFIED